



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ered with the 12-inch telescope. But measures of such pairs have not been made with this instrument.

It would have been an easy matter to have increased the number of discoveries considerably by including pairs having distances somewhat greater than 5". This, however, has not seemed desirable, and only a few such pairs have been measured, and the results for one only published. W. J. HUSSEY.

September 18, 1900.

ELLIPTIC ELEMENTS OF COMET 1896 V, GIACOBINI.

In November, 1896, I computed elements of the orbit of this comet from places corresponding to the dates 1896, September 8th, October 5th and 29th. The elements then obtained did not accurately represent the observed path of the comet during the latter part of its apparition. About a year ago I found the origin of the discrepancy in a slightly erroneous position of the comparison-star used for the last observation, and at that time computed the system of elements given below. These elements are based on normal places having the dates 1896, September 8th, October 29th, and 1897, January 4th. The interval included between the extreme dates is only four days less than the apparition period of the comet.

Epoch 1896 November 1.5 G. M. T.

$$M = 0^{\circ} \ 39' \ 56''.5$$

$$\Omega = 193 \ 28 \ 47.4$$

$$i = 11 \ 21 \ 14.7 \left. \vphantom{\begin{matrix} \Omega \\ i \end{matrix}} \right\} 1896.0$$

$$\omega = 140 \ 33 \ 43.4$$

$$\phi = 35 \ 42 \ 7.6$$

$$\log e = 9.766094$$

$$\log a = 0.542898$$

$$\mu = 544''.0765$$

$$\text{Period} = 6.52 \text{ years.}$$

W. J. HUSSEY.

PROPER MOTION OF $\text{O}\Sigma \ 165, 45 \text{ GEMINORUM.}$

OTTO STRUVE discovered a faint companion of *45 Geminorum* about 1842, and first measured its position in 1847. Misled by erroneous observations made in 1877, he thought that the change in the relative positions of the two stars was due to orbital motion. This, however, is not the case. It results entirely from the proper motion of the principal star. This is clearly estab-